
Pressure Effect on Superconducting Critical Temperature According to String Model

Ibrahim Adam Ibrahim Hammad², Mubarak Dirar¹, Nadia Omar Alatta², Rasha Abd Alhai Taha³, Kh. M. Haroun², Rawia A. Elgani³

¹Department of Physics, Faculty of Science, International University of Africa, Khartoum, Sudan

²Department of Physics, Faculty of Education, Alzaiem Alazhari University, Omdurman, Sudan

³Department of Physics, Faculty of Science, Sudan University of Science & Technology, Khartoum, Sudan

Email address:

kshiky1986@gmail.com (I. A. I. Hammad)

To cite this article:

Ibrahim Adam Ibrahim Hammad, Mubarak Dirar, Nadia Omar Alatta, Rasha Abd Alhai Taha, Kh. M. Haroun, Rawia A. Elgani. Pressure Effect on Superconducting Critical Temperature According to String Model. *International Journal of Fluid Mechanics & Thermal Sciences*. Vol. 3, No. 6, 2017, pp. 70-74. doi: 10.11648/j.ijfmnts.20170306.12

Received: March 2, 2017; **Accepted:** March 31, 2017; **Published:** November 28, 2017

Abstract: Superconductivity is generally regarded as one of the most striking and widely used physical phenomenon. Physicists in response have shown sheer interest in scrutinizing superconductivity and constructing theoretical models to explain it. The majority of models derived in this regard neglected some aspects of superconductivity. The link between critical temperature and pressure remains a highly neglected and potentially representing a research gap in this area. Thus, this motivates the researchers to construct a new model on the relation between pressure and superconductors critical temperature using a string model. The study mainly aims to construct theoretical model based on string model in attempt to understand the effect of pressure on critical temperature and superconducting resistance. The results of study reveal that using plasma equation for mechanical and thermal pressure the frequency is obtained. It also finds that treating electrons as string the energy is found in terms temperature and pressure. Further, when the superconducting resistance vanishes the corresponding critical temperature was found. Furthermore, the increasing mechanical pressure increases the critical temperature.

Keywords: Plasma Equation, Superconductivity, Resistivity, Critical Temperature, Pressure

1. Introduction

Superconductivity (S_c henceforth) was first discovered in 1911 by Kamerlingh Onnes. Onnes discovered that the element of mercury exhibited zero resistance at critical temperature ($T_c = 4.1$ K) [1, 2], the conditions in which a superconductor retains its superconducting properties is defined by its critical values. These critical values are critical temperature T_c , the critical magnetic field B_c and the critical current density J_c and depend on the superconducting material [3]. At critical temperature the specimen undergoes a phase transition from a state normal electrical resistivity to superconducting state when the resistance vanishes abruptly below the T_c [4]. Another significant discovery was made by Meissner and Ochsenfeld in 1933 when they realized that the magnetic field is expelled out of the body of the superconductor. Meissner and Ochsenfeld found that the magnetic field penetrates the material at a small distance,

with an order (30–60) nm in metal superconductors. They called it London's penetration depth [5]. Superconductors divide into two classes according to behavior in a magnetic field. All pure samples of superconducting elements, except Nb, exhibit type-1 behaviour and their superconductivity destroyed by a modest applied magnetic field B_c , known as the critical field. The behavior of type-1 and superconductors, at a given temperature T and in a uniform external magnetic field H , can be described as follows: if H is smaller than a critical value $H_c(T)$, the superconductor completely expels the magnetic flux from its interior (Meissner effect); as the external field is increased above the critical value $H_c(T)$, the entire specimen reverts from the superconducting to the normal state [6]. Although Nb is the only element that is type-2 in its pure state, other elements generally become type-2 when the electron mean free path is reduced sufficiently by alloying [7]. In type-2 superconductors, the transition to a normal state is quite

gradual. Where superconductivity is only partially destroyed for $H_{C1} \leq H \leq H_{C2}$. The region between H_{C1} and H_{C2} is known as intermediate state as it contains partially both normal and the superconducting states. H_{C1} is called the lower critical field, whereas H_{C2} is known as upper critical field. At H_{C1} the field begins to penetrate the sample, and the penetration increases until H_{C2} is reached. At H_{C2} , the magnetization vanishes and the sample reaches the normal. Type-2 superconductors exhibit imperfect diamagnetism [8]. Type-2 superconductors are known in which H_{C2} is large as $2.8 \times 10^7 \text{ A/m}$ at absolute zero. Such materials are used now for practical superconducting magnet coils, and are anticipated for use in the generation and distribution of electrical power [9].

In 1957 Bardeen, Cooper, and Schrieffer (BCS) proposed a microscopic theory. This marked the first endeavors in the theoretical understanding of the nature of Sc. The present day theoretical state owes much to this development. They showed that bound electron pairs termed as Cooper pairs carry the supercurrent. There is also an energy gap between the normal and superconducting state [10]. The BCS theory injected two powerful ideas into the collective consciousness: pairing and spontaneous symmetry breaking. Pairing was an essentially new idea, introduced by Cooper and brought to fruition by BCS. The symmetry breaking aspect was mostly implicit in the original BCS work, and in earlier of Fritz London and Landau-Ginzburg; but the depth and success of the BCS theory seized the imagination of the theoretical physics community, and catalyzed and intellectual ferment [11]. The development of Sc lead to discovery of new material of high temperature superconductors (HTSCs), with their superconducting T_c exceeding 23°K , a superconductor usually referred to a HTSC if the T_c exceeds 90°K [12, 13]. The observation of HTSCs in complex layered copper oxides (cuprates) by Bednorz and Muller in 1986 should undoubtedly be rated as one of the greatest experimental discoveries of the last century, whereas identifying and understanding the microscopic origin of high-temperature superconductivity stands as one of the greatest theoretical challenges of this century, they discovered the onset of possible superconductivity at exceptionally high temperatures in a black ceramic material comprising four elements: lanthanum, barium, copper and oxygen. Within the next decade many more complex cuprates were synthesized including the mercury cuprate compounds which, to date, have the highest confirmed critical temperature for a superconducting transition, some $T_c = 135^\circ \text{K}$ at room pressure and approximately 160°K under high applied pressure. The new phenomenon initiated by Bednorz and Muller broke all constraints on the maximum T_c predicted by the conventional theory of low-temperature superconducting metals and their alloys. These discoveries could undoubtedly result in large-scale commercial applications for cheap and efficient electricity production, provided long lengths of superconducting wires operating above the liquid nitrogen temperature (80°K) can be

routinely manufactured [14]. Pressure was very effective in inducing Sc in elements and compounds that are not superconducting at ambient pressure, some of these elements such as Si, Ge and cuprate are shown to be effected by applying pressure, by application of pressure on these superconducting cuprates increases T_c for underdoped and optimally doped compounds. For example T_c of $\text{HgBa}_2\text{Ca}_2\text{Cu}_3\text{O}_{8+\delta}$ (135°K at ambient pressure), which is the highest found among any superconductor, increases up to 164°K at 30 GPa. It is generally considered that application of pressure promotes transfer of charge from reservoir block to superconducting block. In under doped superconductors, the pressure raises T_c by doping while a decrease of T_c is observed in overdoped cuprates. In mercury cooperates such as $\text{HgBa}_2\text{Ca}_2\text{Cu}_3\text{O}_{8+\delta}$ even overdoped samples show an increase of T_c under pressure because intrinsic effects are supposed to be more efficient than charge transfer. A possible explanation is shown that under the action of the pressure, the reservoir blocks shrink more than the superconducting blocks, so that T_c increases due to an enhanced proximity effect. In this respect, when the pressure increase the T_c increases [15, 16, 17, 18]. A plethora of studies investigated the effect of pressure on the T_c using an experimental approach. One of these studies dealt with the pressure dependence of the superconducting critical temperature of the $\text{Tl}_{0.5}\text{Pb}_{0.5}\text{Sr}_2\text{Ca}_{1-x}\text{Y}_x\text{Cu}_2\text{O}_7$ system. The results of this study indicated that the pressure is proportional with T_c [19]. Another study examined the effect of pressure on the properties of molybdenum sulfides comparing the superconducting and the normal states. This paper found that when the pressure increases the T_c decreases [20].

2. Pressure Effect on Superconducting Critical Temperature According to String Model

Plasma equation describes ionized particles in a gaseous or liquid form. This equation describes the electron easy motion. The electrons here behave as ionized particles inside matter.

The plasma medium has two types of pressure; thermal pressure and mechanical pressure (p_m) per particle [21].

Thermal pressure is given by

$$P = n \gamma k T \quad (1)$$

Where, n , γ , k and T ; are the number of particles, friction coefficient, Boltzmann constant and temperature respectively. By introducing the two types of pressure, one finds the equation of motion of plasma particles to be

$$mn \frac{dv}{dx} \frac{dx}{dt} = -\frac{\partial p}{\partial x} + \frac{\partial np_m}{\partial x} + \frac{\partial nV_o}{\partial x} \quad (2)$$

Where, V_o is the crystal field.

Thus

$$\frac{n}{2}mv^2 = -n\gamma kT + np_m + V_o n \quad (3)$$

For harmonic oscillator string particles the displacement is given by

$$x = x_o e^{i\omega t}, \quad v = i\omega x \quad (4)$$

Thus inserting equation (4) in (3) yields

$$-\frac{nm\omega^2 x^2}{2} = -n\gamma kT + np_m + V_o n \quad (5)$$

The positive sign of p_m is due to the fact that the pressure is exerted on the system.

Then, from equation (5) one finds the frequency to be

$$\omega = \frac{1}{x} \sqrt{\frac{2}{m} \sqrt{\gamma kT - V_o - p_m}} \quad (6)$$

Consider the electron as string harmonic oscillator, thus, depends on the quantum mechanic the energy of harmonic oscillator is given by

$$E = \hbar\omega \quad (7)$$

But for harmonic oscillator from (4):

$$T = \frac{1}{2}mv^2 = \frac{-mw^2 x^2}{2} = V \text{ i.e } T = V.$$

Where, T is the kinetic energy and V is the potential energy.

Hence the total energy of harmonic oscillator is

$$E = \hbar\omega = T + V = 2V \quad (8)$$

$$V = eV_e \quad (9)$$

Where, e and V_e are the electronic potential and electron charge respectively.

According to classical laws the resistance is given by

$$R = \frac{V_e}{I} = \frac{V}{eI} = \frac{E}{2eI} = \frac{\hbar\omega}{2eI} \quad (10)$$

Sub equation (6) in (10) yields

$$R = \frac{\hbar}{2eIx} \sqrt{\frac{2}{m} \sqrt{\gamma kT - V_o - p_m}} \quad (11)$$

Splitting R to real part R_s and imaginary part R_i

$$R = R_r + iR_i \quad (12)$$

The $R_s = R_r$

When

$$\gamma kT < p_m - V_o \quad (13)$$

$$T < \frac{p_m - V_o}{\gamma k} \quad (14)$$

Then

$$T_c = \frac{p_m - V_o}{\gamma k} \quad (15)$$

R = imaginary = iR_i , $R_r = 0$, $R_s = 0$

Eq. (15) shows that when the pressure increases the T_c is increases.

Driving both sides of Eq. (3) yields

$$d\left(\frac{1}{2}nmv^2\right) = -dn\gamma kT + dnp_m + dV_o n \quad (16)$$

Integrating both side yields

$$\frac{1}{2}nmv^2 = -n\gamma kT + np_m + V_o n + C_o \quad (17)$$

Or

$$n \left[\frac{1}{2}mv^2 + \gamma kT - p_m - V_o \right] = C_o = \text{const} \quad (18)$$

$$E = \left[\frac{1}{2}mv^2 + \gamma kT - p_m - V_o \right] \quad (19)$$

For harmonic oscillator Eq. (7) and Eq. (19) one finds

$$T = \frac{1}{2}mv^2 = V \quad (20)$$

$$E = 2V + \gamma kT - p_m - V_o \quad (21)$$

Or

$$\hbar\omega = E = 2V + \gamma kT - p_m - V_o \quad (22)$$

$$R = \frac{\langle H \rangle}{eI} = \frac{\gamma kT - p_m + 2V - V_o}{eI} \quad (23)$$

The resistance R can split into two parts, real part R_+ and imaginary part R_- . [22].

$$R = R_+ + R_- \quad (24)$$

Hence

$$R_s = R_+, \quad R = R_- \quad (25)$$

When

$$\gamma kT - p_m + 2V - V_o < 0$$

$$T < \frac{p_m - 2V + V_o}{\gamma k} \quad (26)$$

The R_s resistance vanishes as indicated in equations (23), (24) and (26).

Thus critical temperature is given by

$$T_c = \frac{p_m - 2V + V_o}{\gamma k} \quad (27)$$

Consider the definition of R in terms of potential V, by considering the total energy E to be of the harmonic oscillator then

$$E = 2V = \gamma k T - p_m \quad (28)$$

$$V = \frac{\gamma k T - p_m}{2} \quad (29)$$

$$R = \frac{V}{eI} = \frac{\gamma k T - p_m}{2eI} = R_+ + R_- \quad (30)$$

Then, $R = R_+$

When

$$\gamma k T - p_m < 0 \quad (31)$$

Or

$$T < \frac{p_m}{\gamma k} \quad (32)$$

Thus the critical temperature be

$$T_c = \frac{p_m}{\gamma k} \quad (33)$$

For repulsive electronic force, magnetic repulsive potential force V_m and thermal force exerted by surrounding, in addition to the battery potential force V_o . Thus according to harmonic oscillator model

$$\frac{1}{2} m v^2 = V = V_o + V_m \quad (34)$$

$$E = -\gamma k T - p_m + 2V_o + 2V_m \quad (35)$$

$$V_o = \frac{E + \gamma k T + p_m - 2V_m}{2} \quad (36)$$

$$R = \frac{V_o}{eI} = \frac{E + \gamma k T + p_m - 2V_m}{2eI} = R_+ + R_- \quad (37)$$

Then $R = R_+$

When

$$E + \gamma k T + p_m - 2V_m < 0 \quad (38)$$

Then

$$T < \frac{2V_m - E - p_m}{\gamma k} \quad (39)$$

Thus the critical temperature be

$$T_c = \frac{2V_m - E - p_m}{\gamma k} \quad (40)$$

3. Discussion

By treating electrons as a harmonic string on the one hand [see Equations (7, 8)] and by using quantum resistance on the other; a useful expression of resistance depending on V and I is found [see Eq. (10)]. Based on the assumption that the superconductor resistance is equal to the real part, the equation (14) and (15) show it vanishes below a certain critical temperature. The equation (15) proved that the critical temperature increases as mechanical pressure increases. The result of this equation agrees with some experimental studies [19]. The resulted proportional relation of critical temperature and mechanical pressure is only valid in the case of the cuprate superconductor component [23, 24]. Further, the expression (15) coincides with the conventional superconductors for low pressure and low T_c . The quantum resistance can be R defined in terms of energy instead of potential as equation (23) shows. The resistance here is assumed to be consisting of real R_c and imaginary part. In this case, T_c also increases as pressure increase. The resistance vanishes below T_c according to equations (23, 24, and 27). Defining R in terms of attractive potential [see equation (27)] or repulsive electronic force beside attractive magnetic force, [see equation (37)], and splitting R again to real R_c and negative parts, the critical temperatures T_c [equations (33) and (40)] again depends on pressure, equation (40) shows that when pressure increase the T_c decrease and this agrees with some superconductors compounds. Interestingly this agrees with some experimental studies [20, 25]. However, this result is not desirable because changing material from normal state to superconductors becomes extremely difficult.

4. Conclusion

The plasma equation can give useful expression for energy both for the thermal and mechanical pressure. This expression, together with definition of resistance classically and quantum mechanically lead to critical temperature dependent on mechanical pressure. The uniqueness of this model is emerges from its new approach that treats charge carriers as oscillating strings obeying plasma equation coupled with plank quantum law. It shows that the T_c changes with pressure, in agreement with experiment. However, this model needs promotion to give an exact empirical relation between pressure and critical temperature, besides explaining other HTS phenomena and thus, could represent the standing point for further research in this regard.

References

- [1] Berk, N. and Schrieffer, J. (1966) in Superconductivity. Phys. Rev. Lett, 17, 433.
- [2] Poole, P. and JR. (2000) Handbook of superconductivity, A Harcourt Science and Technology Company, USA.

- [3] Christian Barth, (2013) High Temperature Superconductor Cable Concepts for Fusion Magnets, KIT scientific publishing, ISSN: 1869-1765.
- [4] Charles Kittel, (2005) Introduction to solid state physics, eight editions, library of congress cataloging, USA.
- [5] Sharma, R. (2015) Superconductivity Basics and Applications to Magnets, National Physical Laboratory, India.
- [6] Hook, R. and Hall, E. (2010) Solid state physics, Second addition, library of congress cataloging, USA.
- [7] Michel cyrot and Davor Pavuna. (1992) Introduction to superconductivity and high- T_c materials, world scientific, Singapore.
- [8] Giuseppe Grosso and Giuseppe pastori parravcini. (2003) Solid state physics, second printing, British Library Cataloguing, UK.
- [9] Gupta, C. (2009) Solid state physics second revised enlarged edition, Vikas Publishing House (P) Ltd. UBS Publisher's Distributors, New Delhi.
- [10] Cardwell, A. and Ginley, S. (2003) Handbook of superconducting materials volume 1: superconductivity, materials and processes, Institute of Physics Publishing, UK.
- [11] Neeraj Mehta. (2009) Textbook of engineering physics part II, PHI learning private Limited, New Delhi.
- [12] Uchida. (2015) High temperature superconductivity, the road to higher critical temperature, DOI 10.1007/978-4-431-55300-7.
- [13] Wesche. (2015) Physical properties of high-temperature superconductors, John Wiley & Sons, United Kingdom.
- [14] Alexandrov, S. (2003) Theory of Superconductivity from Weak to Strong Coupling, IOP Publishing Ltd, Bristol and Philadelphia.
- [15] Sin, A., Odier, P., Regueiro, M., Ordando, M., and Cunha, A. (1999) pressure effects in $Hg_{0.82}Re_{0.18}Ba_{(2-y)}Sr_yCa_2Cu_3O_{8+\delta}$, in Xavier Obradors, F. Sandiumenge, J. Fontcuberta (eds.), Applied Superconductivity 1999, volume 1. Large scale applications, proceedings of EUCAS, the fourth European conference on applied superconductivity, held in Sitges, Spain.
- [16] Narlikar, A. (2005) Frontiers in superconducting materials, Springer, Germany.
- [17] Mourachkine. (2004) Room-temperature superconductivity, cambridge international science publishing, UK.
- [18] Angilella and Pucci. (2000) Pressure effect in high- T_c superconductors with n inequivalent layers.
- [19] Jover, D., Wilhelm, H., and Wijngaarden, R. (1997) Pressure dependence of the superconducting critical temperature of the $Tl_{0.5}Pb_{0.5}Sr_2Ca_{1-x}Y_xCu_2O_7$ system, Physical review b, volume 55.
- [20] Alekseevsk, N., Dobrovol'sk, N., Nizhankovsk, V., and Tsebro, V. (1975) Effect of pressure on the properties of molybdenum sulfides in the superconducting and normal states, Institute of Physical Problems, USSR Academy of Sciences, Zh. Eksp. Teor. Fiz. 69, 662-665.
- [21] Altambori, A., Zakaria, A., Dirar, M., ELhussien, A., AbdALgani, R., and Abdalla, A. (2016) Quantum relation between superconductivity resistance and energy gap, global journal of engineering science and researches.
- [22] Zakaria, A., Altambori, A., Dirar, M., ELhussien, A., AbdALgani, R., and Abdalla, A. (2016) Quantum effect of magnetic field in destroying superconductivity, international journal of engineering sciences & management.
- [23] Nikolay Plakida. (2013) High-temperature cuprate superconductors, experiment, theory, and applications, DOI 10.1007/978-3-642-12633-8.
- [24] Jover, D., Wilhelm, H., and Wijngaarden, R. (1996) Pressure dependence of the superconducting critical temperature of the $Tl_{0.5}Pb_{0.5}Sr_2Ca_{1-2x}Y_xCu_{2.7}$ system, physical review, volume 55.
- [25] Parinov, I. (2007) Microstructure and Properties of High-Temperature Superconductors, Springer Berlin Heidelberg New York.